

Product:	Anti-Methamphetamine Antibody Not available						
Synonyms:							
Immunogen:	Methamphetamine-BTG						
	1 mg	PAS10395	Host Species:	Ovis aries (Sheep)	Isotype:	IgG	
Pack sizes: Cat. No.	10 mg	PAS10396	Host Breed:	Texel	Format:	Clear liquid	
			Clonality:	Polyclonal	pH:	7.4	
IgG Concentration:			Buffer:	Buffer:			
Lot dependent. Determined @ 280nm.			20mM Phosphate	20mM Phosphate, 150mM Sodium Chloride.			
Recommended Working Concentration*:			Preservative:	Preservative:			
4.05/			0.09% Sodium A	0.09% Sodium Azide.			
1.25 µg/mL *The application notes include recommended starting dilutions; optimal							
dilutions/concentrations sl	hould be detern	nined by the end user.					
Assessment Method:			Species Reactivity:				
Competitive ELISA.			N/A	N/A			
Method of Purification:		Storage:					
Salt fractionation.			For long term sto Avoid repeated fr	Can be stored for up to 3 months at +2°C - +8°C. For long term storage, aliquot and store at ≤-20 °C. Avoid repeated freeze/thaw cycles. Product should be protected from light exposure.			
Recommende	d Applic	ations:	•				
The antibody is	suitable	for the development of i	mmunoassays or imn	nunoaffinity purification	columns.		
Sensitivity:			Target Specifici	Target Specificity*:			
10 ng/mL Methamphetamine produces 82% inhibition in a competitive ELISA employing Methamphetamine polyclonal antibody.			antibody exhibite <8.575%, MDMA (1S,2S)-(+)-Pseu	ELISA microtitre plate based and expressed as % crossreactivity*. The antibody exhibited crossreactivity for the following compounds: Amphetamine <8.575%, MDMA 137.862%, (+/-)-MDA <8.575%, Ephedrine HCl 26.626%, (1S,2S)-(+)-Pseudoephedrine <8.575%, (1R,2S)-(-)-Norephedrine <8.575% and (1S,2S)-(+)-N-Methyl-Pseudoephedrine <8.575%.			
			* Cross reactivity profile may	* Cross reactivity profile may vary with tracer used.			
Related Produ	icts:		•				
Methamphetam	nine-HRF	P, HRP9720, Kit size: 0.5	i mL.				

Notes & Precautions:

Antibody can be affinity purified on request. This product as supplied is intended for research applications only, not for use in therapeutic or diagnostic applications without the expressed written authorization of Randox BioReagents. A safety data sheet (SDS) can be supplied upon request. Vial should be centrifuged briefly before opening to ensure all material is removed from the vial cap.

28-Apr-22 Information correct at time of going to print

